

Unusual usage of wobble modifications in mitochondrial tRNAs of the nematode *Ascaris suum*

Masayuki Sakurai^{a,b}, Takashi Ohtsuki^{a,*}, Tsutomu Suzuki^{a,b}, Kimitsuna Watanabe^{a,*}

^a Department of Integrated Biosciences, Graduate School of Frontier Sciences, The University of Tokyo, Kashiwa, Chiba 277-8562, Japan

^b Department of Chemistry and Biotechnology, Graduate School of Engineering, The University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan

Received 5 April 2005; accepted 7 April 2005

Available online 20 April 2005

Edited by Lev Kisselev

Abstract To understand the decoding property of nematode mitochondrial tRNAs with unusual secondary structures, post-transcriptional modifications at wobble positions of *Ascaris suum* mitochondrial tRNAs corresponding to two-codon families ending with a purine were analyzed. 5-Carboxymethylaminomethyluridine (cmnm⁵U) was identified at the wobble positions of tRNA^{Lys}, tRNA^{Glu} and tRNA^{Gln}, while 5-carboxymethylaminomethyl-2-thiouridine (cmnm⁵s²U) was present in tRNA^{Leu}_{UAA} and tRNA^{Trp}. In most bacterial and mitochondrial tRNAs, the 2-thiouridine derivative is present in tRNAs for Lys, Glu and Gln. These is no report that cmnm⁵s²U is used in tRNA^{Leu}_{UAA} and tRNA^{Trp}. The unusual usage of wobble modifications might assist decoding of nematode mitochondrial mRNAs. © 2005 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: Modified nucleoside; Nematode; Mitochondrial tRNA; T-armless tRNA; Wobble position; 2-Thiouridine

1. Introduction

Post-transcriptional modifications are known to play an important role in tRNA function and structure. Modified nucleosides at the wobble position are required for accurate decoding of genetic information. The wobble position of the tRNA responsible for decoding all four codons in a four-codon family box is mostly occupied by unmodified uridine [1], whereas the wobble uridine of the tRNA corresponding to two-codon families ending with a purine (NNR) is usually modified to discriminate cognate codons from near-cognate

codons [2,3]. Those uridines at the wobble position that are modified with a C5 substituent or its 2-thio derivative have been characterized in detail [4–13].

In animal mitochondrial (mt) tRNAs, unique modifications have been found at the wobble position. Recently, taurine-containing modified uridines (5-taurinomethyluridine, τ m⁵U, and 5-taurinomethyl-2-thiouridine, τ m⁵s²U) were identified in mammalian mt tRNAs corresponding to NNR codons [14]. The lack of these modifications causes certain mitochondrial diseases in humans [15,16]. Such taurine-containing modifications have also been found in urochordate mt tRNA^{Gly}_{UCU} [17] (Kondow et al., unpublished observations). However, in yeast [18] and *Tetrahymena* [2], 5-carboxymethylaminomethyluridine (cmnm⁵U) derivatives are present at the wobble position of mt tRNAs corresponding to the two-codon families. We wondered when during the evolution from single-celled eukaryotes to mammals the 5-carboxymethylaminomethyl group was replaced with the 5-taurinomethyl group in mitochondria, with particular interest in the nematode.

Nematode mt tRNAs have extremely unusual secondary structures in that they lack the T-arm or D-arm [19–22] found in the cloverleaf secondary structures of canonical tRNAs [2]. To understand how these tRNAs function in the nematode mt translation system, a characterization of their post-transcriptional modifications is first of all necessary. To date, 15 tRNAs (13 T-armless tRNAs and 2 D-armless tRNA^{Ser}) have been characterized [23–25] in *A. suum* mitochondria. In a previous study, we reported that all T-armless *A. suum* mt tRNAs examined had 1-methyladenosine at position 9 (m¹A₉) and that this modification was indispensable for their structure, aminoacylation, and EF-Tu-binding [25]. These observations exemplify the importance of post-transcriptional modifications for tRNA structure and function.

Within the context of these previous studies, we attempted in the present study to identify the modifications at wobble positions of nematode mt tRNAs corresponding to NNR codons by isolating and determining the RNA sequences of tRNAs from the nematode *A. suum*. Here we report the characteristic wobble modifications of the *A. suum* tRNAs.

2. Materials and methods

2.1. Purification of individual mt tRNAs from *A. suum*

Ascaris suum were provided by Dr. K. Kita and Dr. Y. Watanabe (Univ. of Tokyo). The crude RNA fraction was extracted from 200 g of *A. suum* as described [23]. Individual mt tRNAs were purified by

*Corresponding authors. Fax: +81 86 251 8219 (T. Ohtsuki); +81 3 5530 2064 (K. Watanabe).

E-mail addresses: ohtsuk@cc.okayama-u.ac.jp (T. Ohtsuki), kwatanab@jbirc.aist.go.jp (K. Watanabe).

¹ Present address: Department of Bioscience and Biotechnology, Okayama University, 3-1-1 Tsushima-naka, Okayama 700-8530, Japan.

² Present address: Biological Information Research Center, National Institute of Advanced Industrial Science and Technology, 2-41-6 Aomi, Koto-ku, Tokyo 135-0064, Japan.

Abbreviations: *A. suum*, *Ascaris suum*; APM, (*N*-acryloylamino)phenyl-mercuric chloride; cmnm⁵U, 5-carboxymethylaminomethyluridine; m¹A, 1-methyladenosine; mt, mitochondrial; NNR codons, two-codon families ending with a purine; τ m⁵U, 5-taurinomethyluridine; τ m⁵s²U, 5-taurinomethyl-2-thiouridine; cmnm⁵s²U, 5-carboxymethylaminomethyl-2-thiouridine; 2D-TLC, two-dimensional thin layer chromatography; U₃₄, uridine at position 34

chaplet column chromatography [14], which was invented for tRNA isolation based on a hybrid selection method using solid-phase DNA probes [26]. The sequences of the probes were as follows:

tRNA^{Leu}_{UAA}, 5'-AGTTGTCCCATATCTTTACGCTTAAAACA-3';
tRNA^{Trp}, 5'-TGAAAACCAAGAGTTTAACTTAACTTAAA-3';
tRNA^{Gln}, 5'-CTATACTACAACCCCTTTACACCAAAAATA-3';
tRNA^{Lys}, 5'-AAAAATCTAACACTTTAACTTAAAGTTAAC-3';
tRNA^{Glu}, 5'-CGAAAAAGAAATATACAAAAAATTTACT-3'.

2.2. Sequencing of mt tRNAs

5'- or 3'-³²P labeled *A. suum* mt tRNAs were purified again by gel electrophoresis and sequenced by the method of Donis-Keller [27] using RNases T₁, U₂, PhyM, and CL₃.

2.3. Two-dimensional thin layer chromatography

The nucleotide sequences of tRNA^{Leu}_{UAA}, tRNA^{Trp}, tRNA^{Gln}, tRNA^{Lys}, and tRNA^{Glu}, including modified nucleotides, were analyzed by two-dimensional thin layer chromatography (2D-TLC) as previously described [28].

2.4. Liquid chromatography–mass spectrometry

For nucleoside analysis, each tRNA was purified by PAGE before nuclease digestion. Nucleosides obtained from digestion of mt tRNAs with nuclease P₁ and bacterial alkaline phosphatase were analyzed by LC/MS as described [14,29]. Chromatography was carried out using an ODS reversed-phase column under the conditions reported in [29]. Oligonucleotides produced by RNase T₁ digestion of each tRNA were also analyzed by LC/MS as described in the literature [30,31]. For chromatography, the hydrolysates were applied directly onto a Supelcosil LC-DB C₁₈ column (300 × 1.0 mm; Supelco), and separated using a 33-minute gradient of 5–80% solvent B (50% methanol) in solvent A (0.4 M 1,1,1,3,3,3-hexafluoro-2-propanol adjusted to pH 7.0 with triethylamine).

2.5. Phenyl-mercuric gel electrophoresis

The presence of the 2-thio derivative in mt tRNAs for Leu(UUR) and Trp was verified by electrophoresis using a 10% polyacrylamide gel containing 7 M urea and 0.015 mg/ml (*N*-acryloylamino)phenyl-mercuric chloride (APM) [32], which was provided by Dr. N. Shigi and Mr. Y. Yamamoto in our laboratory.

3. Results

3.1. Purification and sequencing analysis of mt tRNAs

We isolated mt tRNAs for Leu(UUR), Trp, Gln, Lys, and Glu, all of which read NNR codons. The sequences of these tRNAs were analyzed by the methods of Donis-Keller [27] (Fig. S1) and Kuchino et al. [28] (Fig. S2). We also analyzed nucleosides obtained by nuclease P₁ digestion of each tRNA by LC/MS (Fig. 1 and S3). These analyses resulted in the tRNA sequences shown in Fig. 3. Modified nucleosides in the wobble position of these five tRNAs were compared with other organisms as shown in Table 1.

3.2. Wobble position corresponding to NNR codons

All of the tRNAs which read NNR codons had cmnm⁵U or its 2-thio derivative at the wobble position, but the tm⁵U derivative that has been found in mammalian mt tRNAs [14] was not detected. Unexpectedly, cmnm⁵s²U₃₄ was detected in tRNA^{Leu}_{UAA} (Fig. 1A) and tRNA^{Trp} (Fig. S3[A]) but not in tRNAs for Lys, Gln, or Glu, which have cmnm⁵U₃₄. The presence of these modifications was also confirmed by LC/MS analysis of anticodon-containing fragments obtained by RNase T₁-digested tRNAs for Leu(UUR) and Lys (Fig. 1C–F). As for tRNA^{Leu}_{UAA} and tRNA^{Trp}, not only cmnm⁵s²U₃₄ but also cmnm⁵U₃₄ was detected by LC/MS analysis (Fig. 1A, C and E for tRNA^{Leu}_{UAA}; Fig. S3[A] for tRNA^{Trp}), indicat-

ing that these are partial modifications. On the basis of the UV chromatogram (Fig. 1 and S3) and the enzymatic digestion pattern obtained by Donis-Keller's method (Fig. S1), 90% of the cmnm⁵U₃₄ derivatives in tRNA^{Leu}_{UAA} were found to be the 2-thio modification variant. On the other hand, tRNA^{Trp} consists of three different species that contain 35% cmnm⁵U, 15% cmnm⁵s²U, and 50% unmodified uridine, respectively.

3.3. Phenyl-mercuric gel electrophoresis

The presence of cmnm⁵s²U was further confirmed by electrophoresis in gels containing APM [32]. In this analysis, the mercuric compound in the gel interacts specifically with tRNAs containing a thiocarbonyl group, which retards the migration of these tRNAs. Specific retardation of tRNA^{Leu}_{UAA} and tRNA^{Trp} on the APM gel (Fig. 2) indicated the presence of a thiocarbonyl group in these tRNAs.

4. Discussion

In this study, we analyzed wobble modifications of tRNAs corresponding to NNR codons to evaluate when the tm⁵ group replaced the cmnm⁵ group during the evolution of single-celled eukaryotes to mammals. We found that the cmnm⁵ group is used in nematode mt tRNAs, and that some of them possess a characteristic 2-thio modification of the wobble uridine.

The cmnm⁵ group in the wobble uridine of mt tRNAs may have been replaced with tm⁵ group during evolution, as shown in Table 1. Apart from the 2-thio modification or 2'-*O*-methylation, cmnm⁵U₃₄ derivatives exist in eubacteria; mitochondria of unicellular eukaryotes, such as yeast and *Tetrahymena* (cmnm⁵U₃₄ in tRNA^{Trp}) [2], and nematode mitochondria, while taurine-containing modifications at U₃₄ exist in mitochondria of higher animals such as vertebrates [14] and urochordates [17] (Kondow et al., unpublished observations). Thus, the evolutionary boundary between cmnm⁵U₃₄ and tm⁵U₃₄ use in mt tRNAs is most likely between nematodes and urochordates.

2-Thiouridine (s²U₃₄) derivatives at the anticodon wobble position have been reported in tRNAs for Gln, Lys, and Glu from various sources, including eubacteria and eukaryotic cytoplasm, mitochondria, and chloroplasts, but have never been found in tRNAs for Leu(UUR) and Trp [2,3] (Table 1). In the present study of *A. suum* mt tRNAs, an s²U₃₄ derivative was found in tRNA^{Leu}_{UAA} and tRNA^{Trp}. In contrast, there was no s²U derivative in *A. suum* mt tRNA^{Gln}, tRNA^{Lys}, or tRNA^{Glu}, although in eubacteria these tRNA species usually have an s²U₃₄ derivative [2,3]. Why do nematode mt tRNAs use this s²U₃₄ derivative? It is possible that the 2-thio group of the wobble uridine in *A. suum* mt tRNAs is necessary to raise the efficiency of translating NNR codons. UUR codons appear quite frequently in nematode mt protein genes, and have been found to comprise 12–13% of all codons [20,33]. This value is much higher than the average usage of other two-codon families (about 3%). Thus, translation errors involving UUR codons would result in serious consequences for mt protein synthesis. In addition, UUY codons specifying phenylalanine also occur frequently (13–15%) [20], so that it is very important for the nematode mt translation system to avoid the misreading of tRNA^{Leu}_{UAA} against UUY codons. The 2-thio modification leads to an increase in the 3'-endo conformation of the




Fig. 1. LC/MS analysis of *A. suum* mt tRNA^{Leu}_{UAA} and tRNA^{Lys}. (A, B) Total nucleoside analysis of purified *A. suum* mt tRNA^{Leu}_{UAA} and tRNA^{Lys}, respectively. (top) UV chromatograms for nucleosides. (bottom) Mass chromatograms for cmnm⁵s²U and cmnm⁵U with mass filters at *m/z* 332 and 348. The relative peak intensity of cmnm⁵s²U and cmnm⁵U in the mass chromatograms are adjusted against that of m¹A as the standard (100%). (C, D) Chromatograms for RNase T₁-digested tRNA^{Leu}_{UAA} and tRNA^{Lys}, respectively. (top) UV traces at 260 nm; numbers indicate elution positions of RNA fragments as follows: (C) 1, UGp; 2, UUGp; 3, AAGp; 4, UUAUm¹AGp; 5, AUAUGp; 6, CAΨΨUGp; 7, CAUAAAGp; 8, ΨAAAAAGp; 9, ACAACUCCA; 10, ΨΨUcmnm⁵s²UAAm¹GCGp. (D) 1, CCA; 2, UGp; 3, UUGp; 4, AAAUGp; 5, Ψm²GΨΨAGp; 6, UUUAAAGp; 7, UUm¹ACUAAAGp; 8, AUUcmnm⁵UUUt⁶AAΨCUGp. (middle) Base peak mass chromatograms. (bottom) Mass chromatograms for *m/z* 1505.9 (tRNA^{Leu}_{UAA}) and *m/z* 1342.6 (tRNA^{Lys}) to detect doubly charged and triply charged ions of anticodon-containing fragments, respectively. (E, F) Deconvoluted mass spectra for anticodon-containing fragments derived from tRNA^{Leu}_{UAA} and tRNA^{Lys}, respectively. The calculated *m/z* values are given for each peak. The peaks of fragments with 3'-cyclic ends are shown with asterisks.

uridine nucleotide [4,34]. Thus, the 2-thio modification of U₃₄ has been considered to be involved in the rigidity of the anticodon structure [4–7], which restricts the tRNA to recognizing

NNR codons and not NNY codons [11,12]. Furthermore, this modification has been shown to contribute to aminoacylation efficiency [8,9] and the translation rate [10,13]. The 2-thio




Fig. 2. Phenyl-mercuric gel electrophoresis of mt tRNAs to detect specific retardation due to the thiocarbonyl group. 5'-Labeled tRNAs were analyzed on denaturing 10% polyacrylamide gels with (left, APM+) or without (right, APM-) APM.




Fig. 3. Nucleotide sequence and secondary structure of *A. suum* mt tRNA^{Leu_{UAA}}, tRNA^{Trp}, tRNA^{Gln}, tRNA^{Lys}, and tRNA^{Glu}. The letters in the parentheses indicate partial modification or genetic polymorphism.

Table 1
Modified nucleosides in the wobble position of tRNA

	UYR (Y = U or C, R = A or G)		VAR (V = C, A or G, R = A or G)		
	Leu(UUR)	Trp	Gln	Lys	Glu
Nematode mt <i>A. suum</i>	cmnm ⁵ s ² U, cmnm ⁵ U	U, cmnm ⁵ U, cmnm ⁵ s ² U	cmnm ⁵ U	cmnm ⁵ U	cmnm ⁵ U
Eubacteria <i>E. coli</i> <i>M. capricolum</i>	cmnm ⁵ Um [35] cmnm ⁵ U	(inexistent) cmnm ⁵ Um	s ² U* [36] cmnm ⁵ U	mm ⁵ s ² U cmnm ⁵ U	nm ⁵ s ² U cmnm ⁵ U
Fungous mt <i>S. cerevisiae</i>	cmnm ⁵ U [18]	cmnm ⁵ U [18]		cmnm ⁵ U* [18]	
Vertebrate mt Bovine Human	τm ⁵ U [14] τm ⁵ U [14]	U* U*		τm ⁵ s ² U [14] τm ⁵ s ² U [14]	U* U*

Blanks indicate unidentified nucleosides. Numbers in parentheses denote references, and modified nucleotides without parenthesis are referred to <http://www.uni-bayreuth.de/departments/biochemie/trna/> [2] except for those of *A. suum* mt tRNAs. Asterisks show unknown derivatives of the indicated modified nucleosides.

group at wobble uridine in *A. suum* mt tRNA^{Leu}_{UAA} probably raises translation efficiency and specificity for UUR codons to prevent the possible errors by discriminating from UUY codons. The other two-codon sets, CAR (Gln), AAR (Lys), GAR (Glu), and UGR (Trp), are not used frequently (1.2–3.2% in mt protein genes of nematode) [20]. This might explain why tRNAs for Gln, Lys, and Glu do not have a 2-thio modification and why tRNA^{Trp} has only a partial 2-thio modification.

In conclusion, we found unusual use of the s²U₃₄ derivative in nematode mt tRNAs, which might raise the translation efficiency of frequently used codons. We also found that nematode mt tRNAs for NNR codons have the cmnm⁵ group at the wobble uridine. The τm⁵ modification of the wobble uridine, which is related to human mitochondrial disease, was generated probably during the course of evolution between nematodes and urochordates. Further analysis of the functional differences between τm⁵ and cmnm⁵ groups and the co-evolution of the ribosome site responsible for codon–anticodon interaction during the change from cmnm⁵ to τm⁵ should shed further light on the biological meaning of this evolution.

Acknowledgments: This work was supported by Grants-in-Aid from the Ministry of Culture, Sports, Science and Technology, Japan to K.W. and T.O. We thank Dr. Y. Watanabe for materials and valuable discussions, and Dr. K. Kita, Dr. N. Shigi and Mr. Y. Yamamoto for materials. We are grateful to Dr. Takeo Suzuki and Mr. Y. Ikeuchi for their technical assistance on tRNA isolation and mass spectrometry.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version at [doi:10.1016/j.febslet.2005.04.009](https://doi.org/10.1016/j.febslet.2005.04.009).

References

[1] Yokobori, S., Suzuki, T. and Watanabe, K. (2001) Genetic code variations in mitochondria: tRNA as a major determinant of genetic code plasticity. *J. Mol. Evol.* 53, 314–326.

[2] Sprinzl, M., Horn, C., Brown, M., Ioudovitch, A. and Steinberg, S. (1998) Compilation of tRNA sequences and sequences of tRNA genes. *Nucleic Acids Res.* 26, 148–153.

[3] Bjork, G.R. (1998) in: *Modification and Editing of RNA* (Grosjean, H. and Benne, R., Eds.), pp. 577–581, ASM Press, Washington, DC.

[4] Yokoyama, S., Watanabe, T., Murao, K., Ishikura, H., Yamaizumi, Z., Nishimura, S. and Miyazawa, T. (1985) Molecular mechanism of codon recognition by tRNA species with modified uridine in the first position of the anticodon. *Proc. Natl. Acad. Sci. USA* 82, 4905–4909.

[5] Yokoyama, S. (1995) in: *TRNA: Structure, Biosynthesis, and Function* (Soll, D. and RajBhandary, U., Eds.), pp. 207–223, American Society for Microbiology, Washington, DC.

[6] Benas, P., Bec, G., Keith, G., Marquet, R., Ehresmann, C., Ehresmann, B. and Dumas, P. (2000) The crystal structure of HIV reverse-transcription primer tRNA^{Lys}(3) shows a canonical anticodon loop. *RNA* 6, 1347–1355.

[7] Sundaram, M., Durant, P.C. and Davis, D.R. (2000) Hypermodified nucleosides in the anticodon of tRNA^{Lys} stabilize a canonical U-turn structure. *Biochemistry* 39, 12575–12584.

[8] Madore, E., Florentz, C., Giege, R., Sekine, S., Yokoyama, S. and Lapointe, J. (1999) Effect of modified nucleotides on *Escherichia coli* tRNA^{Glu} structure and on its aminoacylation by glutamyl-tRNA synthetase. Predominant and distinct roles of the mm⁵ and s² modifications of U34. *Eur. J. Biochem.* 266, 1128–1135.

[9] Kruger, M.K., Pedersen, S., Hagervall, T.G. and Sorensen, M.A. (1998) The modification of the wobble base of tRNA^{Glu} modulates the translation rate of glutamic acid codons in vivo. *J. Mol. Biol.* 284, 621–631.

[10] Kruger, M.K. and Sorensen, M.A. (1998) Aminoacylation of hypomodified tRNA^{Glu} in vivo. *J. Mol. Biol.* 284, 609–620.

[11] Ashraf, S.S., Sochacka, E., Cain, R., Guenther, R., Malkiewicz, A. and Agris, P.F. (1999) Single atom modification (O → S) of tRNA confers ribosome binding. *RNA* 5, 188–194.

[12] Yarian, C., Townsend, H., Czeszkowski, W., Sochacka, E., Malkiewicz, A.J., Guenther, R., Miskiewicz, A. and Agris, P.F. (2002) Accurate translation of the genetic code depends on tRNA modified nucleosides. *J. Biol. Chem.* 277, 16391–16395.

[13] Urbonavicius, J., Qian, Q., Durand, J.M., Hagervall, T.G. and Bjork, G.R. (2001) Improvement of reading frame maintenance is a common function for several tRNA modifications. *EMBO J.* 20, 4863–4873.

[14] Suzuki, T., Suzuki, T., Wada, T., Saigo, K. and Watanabe, K. (2002) Taurine as a constituent of mitochondrial tRNAs: new insights into the functions of taurine and human mitochondrial diseases. *EMBO J.* 21, 6581–6589.

[15] Yasukawa, T., Suzuki, T., Ishii, N., Ohta, S. and Watanabe, K. (2001) Wobble modification defect in tRNA disturbs codon–

- anticodon interaction in a mitochondrial disease. *EMBO J.* 20, 4794–4802.
- [16] Kirino, Y., Yasukawa, T., Ohta, S., Akira, S., Ishihara, K., Watanabe, K. and Suzuki, T. (2004) Codon-specific translational defect caused by a wobble modification deficiency in mutant tRNA from a human mitochondrial disease. *Proc. Natl. Acad. Sci. USA* 101, 15070–15075.
- [17] Kondow, A., Suzuki, T., Yokobori, S., Ueda, T. and Watanabe, K. (1999) An extra tRNA^{Gly} (U^{*}CU) found in ascidian mitochondria responsible for decoding non-universal codons AGA/AGG as glycine. *Nucleic Acids Res.* 27, 2554–2559.
- [18] Martin, R.P., Sibley, A.P., Gehrke, C.W., Kuo, K., Edmonds, C.G., McCloskey, J.A. and Dirheimer, G. (1990) 5-Carboxymethylaminomethyluridine is found in the anticodon of yeast mitochondrial tRNAs recognizing two-codon families ending in a purine. *Nucleic Acids Res.* 29, 956–959.
- [19] Wolstenholme, D.R., Macfarlane, J.L., Okimoto, R., Clary, D.O. and Wahleithner, J.A. (1987) Bizarre tRNAs inferred from DNA sequences of mitochondrial genomes of nematode worms. *Proc. Natl. Acad. Sci. USA* 84, 1324–1328.
- [20] Okimoto, R., Macfarlane, J.L., Clary, D.O. and Wolstenholme, D.R. (1992) The mitochondrial genomes of two nematodes, *Caenorhabditis elegans* and *Ascaris suum*. *Genetics* 130, 471–498.
- [21] Keddie, E.M., Higazi, T. and Unnasch, T.R. (1998) The mitochondrial genome of *Onchocerca volvulus*: sequence, structure and phylogenetic analysis. *Mol. Biochem. Parasitol.* 95, 111–127.
- [22] Hu, M., Chilton, N.B. and Gasser, R.B. (2002) The mitochondrial genomes of the human hookworms, *Ancylostoma duodenale* and *Necator americanus* (Nematoda: Secernentea). *Int. J. Parasitol.* 32, 145–158.
- [23] Watanabe, Y., Tsurui, H., Ueda, T., Furushima, R., Takamiya, S., Kita, K., Nishikawa, K. and Watanabe, K. (1994) Primary and higher order structures of nematode (*Ascaris suum*) mitochondrial tRNAs lacking either the T or D stem. *J. Biol. Chem.* 269, 22902–22906.
- [24] Watanabe, Y., Tsurui, H., Ueda, T., Furusihima-Shimogawara, R., Takamiya, S., Kita, K., Nishikawa, K. and Watanabe, K. (1997) Primary sequence of mitochondrial tRNA(Arg) of a nematode *Ascaris suum*: occurrence of unmodified adenosine at the first position of the anticodon. *Biochim. Biophys. Acta* 1350, 119–122.
- [25] Sakurai, M., Ohtsuki, T. and Watanabe, K. (2005) Modification at position 9 with 1-methyladenosine is crucial for structure and function of nematode mitochondrial tRNAs lacking the entire T-arm. *Nucleic Acids Res.* 33, 1653–1661.
- [26] Wakita, K., Watanabe, Y., Yokogawa, T., Kumazawa, Y., Nakamura, S., Ueda, T., Watanabe, K. and Nishikawa, K. (1994) Higher-order structure of bovine mitochondrial tRNA(Phe) lacking the 'conserved' GG and T psi CG sequences as inferred by enzymatic and chemical probing. *Nucleic Acids Res.* 22, 347–353.
- [27] Donis-Keller, H. (1980) Phy M: an RNase activity specific for U and A residues useful in RNA sequence analysis. *Nucleic Acids Res.* 8, 3133–3142.
- [28] Kuchino, Y., Hanyu, N. and Nishimura, S. (1987) Analysis of modified nucleosides and nucleotide sequence of tRNA. *Methods Enzymol.* 155, 379–396.
- [29] Pomerantz, S.C. and McCloskey, J.A. (1990) Analysis of RNA hydrolyzates by liquid chromatography–mass spectrometry. *Methods Enzymol.* 193, 796–824.
- [30] Kaneko, T., Suzuki, T., Kapushoc, S.T., Rubio, M.A., Ghazvini, J., Watanabe, K. and Simpson, L. (2003) Wobble modification differences and subcellular localization of tRNAs in *Leishmania tarentolae*: implication for tRNA sorting mechanism. *EMBO J.* 22, 657–667.
- [31] Soma, A., Ikeuchi, Y., Kanemasa, S., Kobayashi, K., Ogasawara, N., Ote, T., Kato, J., Watanabe, K., Sekine, Y. and Suzuki, T. (2003) An RNA-modifying enzyme that governs both the codon and amino acid specificities of isoleucine tRNA. *Mol. Cell* 12, 689–698.
- [32] Igloi, G.L. (1988) Interaction of tRNAs and of phosphorothioate-substituted nucleic acids with an organomercurial. Probing the chemical environment of thiolated residues by affinity electrophoresis. *Biochemistry* 27, 3842–3849.
- [33] Nakamura, Y., Gojobori, T. and Ikemura, T. (2000) Codon usage tabulated from international DNA sequence databases: status for the year 2000. *Nucleic Acids Res.* 28, 292.
- [34] Sierzputowski, H., Sochacka, E., Malkiewicz, A., Kuo, K., Gehrke, C.W. and Agris, P.F. (1987) Chemistry and structure of modified uridines in the anticodon, wobble position of transfer-RNA are determined by thiolation. *J. Am. Chem. Soc.* 109, 7171–7177.
- [35] Horie, N., Yamaizumi, Z., Kuchino, Y., Takai, K., Goldman, E., Miyazawa, T., Nishimura, S. and Yokoyama, S. (1999) Modified nucleosides in the first positions of the anticodons of tRNA(Leu)4 and tRNA(Leu)5 from *Escherichia coli*. *Biochemistry* 38, 207–217.
- [36] Yaniv, M. and Folk, W.R. (1975) The nucleotide sequences of the two glutamine transfer ribonucleic acids from *Escherichia coli*. *J. Biol. Chem.* 250, 3243–3253.